

Project SANC:

(Support of Analytic and Numeric calculations for experiments at Colliders)

Ideas and Realization

A. Andonov, A. Arbuzov, D. Bardin, S. Bondarenko,

P. Christova, L. Kalinovskaya, G. Nanava

Laboratory of Nuclear Problems, Joint Institute for Nuclear Research

OUTLINE

1. Ideas

2. Basic notions

3. Realization: <http://brg.jinr.ru>, or <http://pcjinr01.cern.ch>

- Decays $B \rightarrow f\bar{f}$
- Decays $H \rightarrow \gamma\gamma (Z\gamma, ZZ, WW), Z \rightarrow WW$
- Processes $2f \rightarrow 2f$ (talk by Lida Kalinovskaya)

4. Demonstration of version 0.21

1. Ideas

- main idea of the project is to create a tool similar to CompHEP, i.e.
 - many processes ME MC generator;
- but:
 - at one-loop precision level (QED, EW, QCD);
 - keeping all external particle masses finite;
 - * whenever necessary (e.g. processes with top-quark);
 - assuming REI (Reactions of Experimental Interest) criterion, i.e.
 - * initial particles (electrons, photons, partons) are massless;
 - * actually fermions of first generation are massless whenever possible;
- present phase assumes to reach a description of any 5 particle processes:
 - 2 → 3 processes;
 - 1 → 4 decays;
 - * we prefer to say 5 → 0

Of course, there is GRACE LOOP version 2.xx and many others...

Network system SANC: technical details

SANC SERVER

- acts as an application provider. It provides access to various applications written in several computer languages
- provides functionality to access database servers, builds SANC-specific database structure and manipulates SANC data
- supports list of its own functions, procedures and methods (Server Functions List – SFL) and exports the list to the other SANC Servers
- provides functionality (via SOAP – PRC) to access and execute the methods from its SFL
- implements Intermediate Access Ideology
- is a user control system
- is an encrypt – decrypt system
- is an internal CVS (Concurrent Versions System)

SANC CLIENT is

- a multi-purpose development environment
- a text editor
- a Database browser
- a project management tool

USED TECHNOLOGIES : JAVA, XML, SOAP

2. Basic notions

- **The Standard Model (SM) Lagrangian in the R_ξ gauge:**

$$\mathcal{L} = \mathcal{L}(\text{IPS of 25 parameters, fields, } \xi_A, \xi_Z, \xi)$$

- **Fields:** fermions, vector bosons, example of Feynman Rules

vector boson propagators:

$$A : \text{~~~~~} \quad \frac{1}{p^2} \left\{ \delta_{\mu\nu} + (\xi_A^2 - 1) \frac{p_\mu p_\nu}{p^2} \right\}$$

$$Z : \text{~~~~~} \quad \frac{1}{p^2 + M_Z^2} \left\{ \delta_{\mu\nu} + (\xi_Z^2 - 1) \frac{p_\mu p_\nu}{p^2 + \xi_Z^2 M_Z^2} \right\}$$

$$W^\pm : \text{~~~~~} \quad \frac{1}{p^2 + M_W^2} \left\{ \delta_{\mu\nu} + (\xi^2 - 1) \frac{p_\mu p_\nu}{p^2 + \xi^2 M_W^2} \right\}$$

- **Unphysical fields:** $\phi^0, \phi^\pm, Y^A, Y^Z, X^\pm$, the physical H field.

- **Passarino–Veltman (PV) functions and reduction:**

Presently, **SANC** knows ALL up to third rank tensorial reduction of up to four-point PV functions: A_0, B_0 , and D_0 ; and the *auxiliary* PV functions: a_0, b_0, c_0 and d_0 , which are due to particular form of photon propagator in R_ξ gauge (terms $\propto 1/p^4$).

- **A fortran library** for numerical calculation of these functions is created and thoroughly tested by means comparison with the other codes.

Amplitude's basis, Scalar Form Factors (**SFF**), Helicity Amplitudes (**HA**)

Example of decays $B(Q) \rightarrow f(p_1)\bar{f}(p_2)$

$$H \rightarrow f\bar{f} \quad - \quad \mathcal{A} \propto I \text{ SFF}_S$$

1 structure (**I**-basis), **1 SFF**, **1 HA**

$$Z \rightarrow f\bar{f} \quad - \quad \mathcal{A} \propto i\gamma_\mu\gamma_6 \text{ SFF}_L + i\gamma_\mu \text{ SFF}_Q + m_f (p_1 - p_2)_\mu \text{ SFF}_D$$

3 structures (**L,Q,D**-basis), **3 SFFs**, **3 HAs**

$$W \rightarrow u\bar{d} \quad - \quad \mathcal{A} \propto i\gamma_\mu\gamma_6 \text{ SFF}_L + i\gamma_\mu\gamma_7 \text{ SFF}_R + m_u (p_1 - p_2)_\mu \gamma_6 \text{ SFF}_{LD} + m_d (p_1 - p_2)_\mu \gamma_7 \text{ SFF}_{RD}$$

4 structures (**L,R,LD,RD**-basis), **4 SFFs**, **4 HAs**

The **3 HAs** depend on kinematical factors, coupling constants and **3 SFFs**, example of Z decay:

$$\mathbf{HA}_{0^{++}}^Z = \mathbf{HA}_{0^{--}}^Z = \frac{gm_f}{c_W} \left[a_f \text{ SFF}_L + \delta_f \text{ SFF}_Q + \frac{1}{2} a_f \beta_f^2 M_Z^2 \text{ SFF}_D \right]$$

$$\mathbf{HA}_{^{++-}}^Z = \frac{gM_Z}{\sqrt{2}c_W} \left[a_f(1 - \beta_f) \text{ SFF}_L + \delta_f \text{ SFF}_Q \right]$$

$$\mathbf{HA}_{^{--+}}^Z = \frac{gM_Z}{\sqrt{2}c_W} \left[a_f(1 + \beta_f) \text{ SFF}_L + \delta_f \text{ SFF}_Q \right]$$

$$\beta_f^2 = 1 - 4\frac{m_f^2}{M_Z^2}, \quad \delta_f = v_f - a_f = -2Q_f s_W^2, \quad a_f = I_f^{(3)}.$$

Amplitude's basis, Scalar Form Factors (**SFF**), Helicity Amplitudes (**HA**), contd.

Example of decays $H(Q) \rightarrow V(p_1)V(p_2)$ [$V = \gamma, Z$]

$$H \rightarrow \gamma\gamma - \mathcal{A} \propto \left(\delta_{\mu\nu} - 2 \frac{p_{1\nu} p_{2\mu}}{M_H^2} \right) \epsilon_\mu^1 \epsilon_\nu^2 \mathbf{SFF}_{\gamma\gamma}, \quad 1 \text{ structure, } \mathbf{1 SFF}, \mathbf{1 HA}$$

$$H \rightarrow Z\gamma - \mathcal{A} \propto \left[\left(1 - \frac{M_Z^2}{M_H^2} \right) \delta_{\mu\nu} - 2 \frac{p_{1\nu} p_{2\mu}}{M_H^2} \right] \epsilon_\mu^1 \epsilon_\nu^2 \mathbf{SFF}_{Z\gamma}, \quad 1 \text{ structure, } \mathbf{1 SFFs}, \mathbf{1 HA}$$

$$H \rightarrow ZZ - \mathcal{A} \propto \left(\delta_{\mu\nu} \mathbf{SFF}_{ZZ}^d + \frac{p_{1\nu} p_{2\mu}}{M_H^2} \mathbf{SFF}_{ZZ}^p \right) \epsilon_\mu^1 \epsilon_\nu^2, \quad 2 \text{ structures, } \mathbf{2 SFFs}, \mathbf{2 HAs}$$

Example of **2 HAs** for $H \rightarrow ZZ$ decay depending on **2 SFFs** (only non-zero **HAs** are shown):

$$\mathbf{HA}_{++}^H = \mathbf{HA}_{--}^H \propto -\mathbf{SFF}_d$$

$$\mathbf{HA}_{00}^H \propto \left[\left(-1 + \frac{1}{2} \frac{M_H^2}{M_Z^2} \right) \mathbf{SFF}_{ZZ}^d + \left(-1 + \frac{1}{4} \frac{M_H^2}{M_Z^2} \right) \mathbf{SFF}_{ZZ}^p \right]$$

From analytic results to numbers

$$d\Gamma(d\sigma) \sim \sum_{\lambda_i \lambda_j \lambda_k \lambda_l} \left| \mathbf{HA}(\mathbf{SFF}^{\mathbf{Born+1-loop}})_{\lambda_i \lambda_j \lambda_k \lambda_l} \right|^2$$


$$d\Gamma(d\sigma) \sim \sum_{\lambda_i \lambda_j \lambda_k \lambda_l} \left| \mathbf{HA}(\mathbf{SFF}^{\mathbf{Born+1-loop+2-loop}})_{\lambda_i \lambda_j \lambda_k \lambda_l} \right|^2$$

A. Ferroglia, G. Passarino, M. Passera, S. Uccirati, Nucl.Phys. B650 (2003) 162-228

+ **Soft bremsstrahlung**

+ **Hard bremsstrahlung either from an analytic calculations**

or from a (partner) MC, e.g. WINHAC



First application of SANC for decays

1. COMPARISON OF SANC WITH KORALZ AND PHOTOS.





By A. Andonov (Dubna, JINR), S. Jadach (Cracow, INP), G. Nanava (Dubna, JINR), Z. Was (Cracow, INP and CERN). CERN-TH-2002-315, Dec 2002.

Published in Acta Phys. Polon. B34 p.2665-2672, 2003.

hep-ph/0212209


2. HOW TO USE SANC TO IMPROVE THE PHOTOS MONTE CARLO SIMULATION OF BREMSSTRAHLUNG IN LEPTONIC W BOSON DECAYS.

By G. Nanava (Dubna, JINR), Z. Was (Cracow, INP and CERN). CERN-TH-2003-075, Mar 2003. Submitted to Acta Phys. Polon. hep-ph/0303206




$\frac{d\Gamma^{1-loop}}{dE_\mu}$

Muon Energy distribution




SANC/KORALZ








$\frac{d\Gamma^{1-loop}}{d\cos\theta_{acol}}$



$\mu^+ \mu^-$ Acolinearity Distribution



SANC/KORALZ









ANNEX

1. Examples of precomputation chains
2. Classification of processes in the SM
3. Examples of some $2 \rightarrow 2$ and $2 \rightarrow 3$ hard processes


Chain: Self \rightarrow Fermion \rightarrow Fermion Self



Two point fermionic diagrams.





Tadpoles: bosonic part,






fermionic part

Chain: Self \rightarrow Boson \rightarrow Boson Self



Two point bosonic diagrams,

fermionic component





One point bosonic diagrams.

Tadpoles: bosonic part,

fermionic part

Chain: Vertex \rightarrow Bff \rightarrow B \rightarrow ff



Vertices: FBF-topology,

BFB-topology.

Classification of processes in the SM

1. PROCESSES $3 \rightarrow 0$

- $Bf\bar{f} \rightarrow 0$ decays

- $H \rightarrow f\bar{f} \longrightarrow 1$ scalar form factor (SFF)
- decay $Z \rightarrow f\bar{f}$ and transition $\gamma^* \rightarrow f\bar{f} \longrightarrow 3$ (SFF)
- decay $W \rightarrow f\bar{f}'$ and crossing-channel $t \rightarrow W^+b \longrightarrow 4$ (SFF)

- $3B \rightarrow 0$ decays and fusions

- decays $H \rightarrow \gamma\gamma, H \rightarrow Z\gamma, H \rightarrow ZZ, W^+W^- \longrightarrow$ correspondingly 1,1,2,2 (SFF)
- fusion $2 \rightarrow 1$
dominant Higgs boson production channel at hadronic colliders $gg \rightarrow H$
- virtual transitions $Z^* \rightarrow W^+W^-$ and $\gamma^* \rightarrow W^+W^-$
“building blocks” for more complicated processes

Decay rates are typical pseudo-observables.

2. PROCESSES $4 \rightarrow 0$

• Processes $4f \rightarrow 0$

- Neutral Current (NC): $f\bar{f} \rightarrow (\gamma, Z, H) \rightarrow f'\bar{f}'$
→ 4 SFF (for massless fermions) & → 10 SFF (for massive fermions)
- Charged Current (CC):
 - * $f_1\bar{f}_2 \rightarrow (W) \rightarrow f_3\bar{f}_4$ → 16 SFF (all fermions are massive)
 - * $F \rightarrow 3f$ — μ, τ decays, $t \rightarrow b\bar{u}$
 - * partonic processes at hadron colliders, for instance: $q_i + \bar{q}_j \rightarrow \bar{b} + t$ ($b + \bar{t}$)

• Processes $2B2f \rightarrow 0$

- Compton effect: $\gamma e \rightarrow \gamma e$
- radiative decays: $Z \rightarrow f\bar{f}\gamma$, $H \rightarrow f\bar{f}\gamma$
- $e^+e^- \rightarrow W^+W^-, ZZ, Z\gamma, \gamma\gamma$
- partonic processes at hadron colliders, for instance: $q_i + \bar{q}_j \rightarrow W(Z) + H$

• Processes $4B \rightarrow 0$

- light-by-light scattering: $\gamma\gamma \rightarrow \gamma\gamma$
- virtual transitions $ZZ \rightarrow ZZ$ (and many others)

3. PROCESSES 5 → 0

- Processes 4fB → 0

- radiative $f\bar{f}\gamma$ production in e^+e^- annihilation: $e^+e^- \rightarrow f\bar{f}\gamma$
- or some partonic processes at LHC:

$$\begin{aligned}g + q_i &\rightarrow q_j + \bar{b} + t, \\q_i + q_i &\rightarrow q_j + q_j + H\end{aligned}$$

- or Higgs decay into four leptons (muons):

$$H \rightarrow \mu^+ + \mu^- + \mu^+ + \mu^-$$

- Processes 3B2f → 0

- one more well-known partonic process at LHC:

$$g + g \rightarrow t + \bar{t} + H$$

– ...

- Processes 5B → 0 ...

4. PROCESSES 6 → 0 ...